

# Suprasorb X – a new Hydrobalance wound dressing and its use in chronic wounds

Eberlein T<sup>1</sup>, Fendler H<sup>2</sup>, Mustafi N<sup>3</sup>, Sauer B<sup>4</sup>, Krenmayer H<sup>5</sup>, Schmitz M<sup>6</sup>

<sup>1</sup> MD, Feucht, Germany

<sup>2</sup> Gesundheitsmanager, Schwaig, Germany

<sup>3</sup> Krankenhaus NordWest, Frankfurt a. Main, Germany

<sup>4</sup> Sauer & Knorr, D-56566 Neuwied

<sup>5</sup> Deutsches Institut für Wundheilung, D-78315 Radolfzell

<sup>6</sup> Lohmann & Rauscher GmbH & Co KG, D-56587 Rengsdorf



Gesundheitsverbund HBH-Kliniken  
Nah bei Ihnen



## Introduction:

The treatment of chronic wounds is a daily challenge for all nurses and physicians. Beside the basic-therapy of the disease, the selection of the adequate wound dressing plays an important role. Regarding to the growing number of materials and products on the market the situation is more and more complex for the prescriber and the user. One approach are products with a broad use in the different stages of exudation and wound healing. Further practical experiences with a biosynthesised cellulose wound dressing, Suprasorb® X, in a open multi-center clinical evaluation study will be presented.

## Material and Methods

The study was conducted on 45 patients with ulcers of different pathogenesis (chronic leg ulcer, diabetic foot ulcer, postsurgical secondary healing wounds and other; wound area 2,8-115 cm<sup>2</sup>, wound age 2 months – 7 years, moderate to medium exudation, period of observation about 4 weeks). The treatment consists of moist wound management: Suprasorb® X and a secondary dressing selected dependent on the level of exudation (film-, foam-, high absorbent dressing).






Fig. 1



## Results:

The results of the study demonstrate the following after 4 weeks: The wound dressings stayed on the wounds up to 7 days depending on the wound situation (averaged 3,82 days, -2,8, +3,18). Significant reduction of wound area (from 5,8 cm<sup>2</sup> to 1,2 cm<sup>2</sup> averaged, Fig. 1, p<0.05) could also be observed. Efficient wound bed cleansing, fundamental growth of granulation tissue and considerably growth of epithelial tissue have also been observed (comparable with first experiences in the USA<sup>1</sup>). Easy handling is a positiv effect for the user, a noticeable pain reduction is one of the benefits for the patient.

## Conclusion

These 45 cases show an excellent clinical efficacy of the hydrobalanced wound dressing, Suprasorb® X, promoting the healing process in all wound healing stages and phases with moderate to medium exudation. Suprasorb® X is a product with a broad use and can reduce the number of products needed in daily use.

## Reference:

1: Alvarez O.M.; Patel M.; Booker J.; Markowitz L.  
Effectiveness of a Biocellulose Wound Dressing for the treatment of Chronic Venous Leg Ulcers, Results of a Single Center Randomized Study Involving 24 Patients; Wounds (2004) 16 (7): 224-233

Conférences Plaies et Cicatrisation (CPC), Paris 14-16. Janv. 2007